arXiv:math/0702760v1 [math.CV] 26 Feb 2007

A CARLESON TYPE CONDITION FOR INTERPOLATING SEQUENCES IN THE HARDY SPACES OF THE BALL OF $\mathbb{C}^n$.

ERIC AMAR

Abstract. In this work we prove that if S is a dual bounded sequence of points in the unit ball $\mathbb{B}$ of $\mathbb{C}^n$ for the Hardy space $H^p(\mathbb{B})$, then S is $H^s(\mathbb{B})$ interpolating with the linear extension property for any $s \in [1, p[$.

1. Introduction.

Let $\mathbb{B}$ be the unit ball of $\mathbb{C}^n$ and σ the Lebesgue's measure on $\partial\mathbb{B}$. As usual we define the Hardy spaces $H^p(\mathbb{B})$ as the closure in $L^p(\partial\mathbb{B})$ of the holomorphic polynomials and $H^\infty(\mathbb{B})$ as the algebra of all bounded holomorphic functions in $\mathbb{B}$.

If $a \in \mathbb{B}$ we note $k_a(z)$ its Cauchy kernel and $k_{a,p}(z)$ its normalized Cauchy kernel in $H^p(\mathbb{B})$:

$$k_a(z) := \frac{1}{(1-\bar{a}\cdot z)^n},\ k_{a,p}(z) := \frac{k_a}{\|k_a\|_p}.$$

We know that $\|k_a\|_p = c(a,p)(1-|a|^2)^{-n/p'}$ with $\frac{1}{p}+\frac{1}{p'} = 1$, $0 < \alpha \le c(a,p) \le \beta$ and α, β independent of $a \in \mathbb{B}$ and of $1 \le p \le \infty$.

We shall need some definitions.

Definition 1.1. *We say that the sequence* $S \subset \mathbb{B}$ *is dual bounded in* $H^p(\mathbb{B})$ *if a dual system* $\{\rho_a\}_{a\in S} \subset H^p(\mathbb{B})$ *for* $\{k_{a,p'}\}_{a\in S}$ *exists and if this sequence is bounded in* $H^p(\mathbb{B})$*, i.e.*

$\exists C > 0 \therefore \forall a \in S,\ \|\rho_a\|_p \le C,\ \forall a, b \in S,\ \langle \rho_a,\ k_{b,p'}\rangle = \delta_{a,b}$.

Definition 1.2. *We say that a sequence* $S \subset \mathbb{B}$ *is* $H^p(\mathbb{B})$ *interpolating for* $1 \le p < \infty$, $S \in IH^p(\mathbb{B})$, *with interpolating constant* $C_I > 0$ *if*

$\forall \lambda \in \ell^p,\ \exists f \in H^p(\mathbb{B}) \therefore \forall a \in S,\ f(a) = \lambda_a \|k_a\|_{p'}$ *and* $\|f\|_p \le C_I \|\lambda\|_p$.

We say that $S \subset \mathbb{B}$ *is* $H^\infty(\mathbb{B})$ *interpolating,* $S \in IH^\infty(\mathbb{B})$, *with interpolating constant* C_I *if*

$\forall \lambda \in \ell^\infty,\ \exists f \in H^\infty(\sigma) \therefore \forall a \in S,\ f(a) = \lambda_a$ *and* $\|f\|_\infty \le C_I \|\lambda\|_\infty$.

Clearly if S is $H^p(\mathbb{B})$ interpolating, then S is dual bounded in $H^p(\mathbb{B})$. In one variable, L. Carleson [4] proved the converse for $p = \infty$ and H. Shapiro & A. Shields [6] did the same for $H^p(\mathbb{D})$, $1 \le p < \infty$.

In [1] we proved, in the general setting of uniform algebras, that if S is H^p interpolating then S is H^s interpolating for any s such that $1 \le s \le p$ provided that some structural hypotheses are verified. This is valid here, in the case of the ball.

Definition 1.3. *We say that the* $H^p(\mathbb{B})$ *interpolating sequence* S *has the linear extension property (LEP) if there is a bounded linear operator* $E\ :\ \ell^p \longrightarrow H^p(\mathbb{B})$ *such that* $\|E\| < \infty$ *and* $\forall \lambda \in \ell^p,\ E\lambda$ *interpolates the sequence* λ *in* $H^p(\mathbb{B})$ *on* S.

In [2] we proved by functional analytic methods still in the general setting of uniform algebras, that if S is dual bounded in H^p then for any s such that $1 \le s \le p$, S is H^s interpolating with the LEP, *provided that* $p = \infty$ *or* $p \le 2$. Hence it remains a gap, the values of $p \in]2, \infty[$.

The aim of this work is to fill this gap in the special case of the ball.

Theorem 1.4. *Let $S \subset \mathbb{B}$ be dual bounded in $H^p(\mathbb{B})$, then S is $H^s(\mathbb{B})$ interpolating with the LEP, for any $s < p$.*

2. Carleson sequences.

Remember that $k_{a,q}$ is the normalized reproducing kernel for the point $a \in \mathbb{B}$ in $H^q(\mathbb{B})$.

Definition 2.1. *We say that the sequence $S \subset \mathbb{B}$ is a Carleson sequence if, for any q such that $1 \leq q < \infty$, we have*

$$\exists D_q > 0,\ \forall \mu \in \ell^q,\ \left\| \sum_{a\in S} \mu_a k_{a,q} \right\|_q \leq D_q \left\| \mu \right\|_q .$$

In fact, up to the duality $H^p(\mathbb{B}) - H^q(\mathbb{B})$, it is proved by Hörmander [5] that this condition, for a $q > 1$, is equivalent to the fact that the measure $\chi := \sum_{a\in S} (1-|a|^2)^n \delta_a$ is a Carleson measure, hence if the condition is true for a $q > 1$ it is true for all q's.

P. Thomas [7] proved that if the sequence S is $H^p(\mathbb{B})$ interpolating for a $p \geq 1$, then γ is a Carleson measure, hence S is a Carleson sequence. As a corollary of this result we have

Lemma 2.2. *If S is dual bounded in $H^p(\mathbb{B})$ for a $p \geq 1$ then S is a Carleson sequence.*

proof:
the dual system $\{\rho_a\}_{a\in S} \subset H^p(\mathbb{B})$ exists and is bounded in $H^p(\mathbb{B})$. Let
$\rho_{a,1} := \rho_a k_{a,p'}$ with p' the conjugate exponent for p. Then we have
$\forall a \in S,\ \left\|\rho_{a,1}\right\|_1 \leq \left\|\rho_a\right\|_p \left\|k_{a,p'}\right\|_{p'} \leq \left\|\rho_a\right\|_p \leq C.$
Moreover $\rho_{a,1}(b) = \rho_a(b)k_{a,p'}(b) = \delta_{ab}\rho_a(a)k_{a,p'}(a) = \delta_{ab}(1-|a|^2)^{-n}$. Hence $\{\rho_{a,1}\}_{a\in S}$ is a dual system for $\{k_{a,\infty}\}_{a\in S}$ and this means that S is also dual bounded in $H^1(\mathbb{B})$. But then it is clear that a dual bounded sequence in $H^1(\mathbb{B})$ is $H^1(\mathbb{B})$ interpolating and we can apply Thomas' theorem to conclude. □

3. The main result.

We are now in position to prove the theorem 1.4.
We shall need the following lemma which was proved for the Poisson Szegö kernel in the ball in [3]. We put on $\partial\mathbb{B}$ the pseudo-distance $\delta(\zeta, z) := \left|1 - \bar{\zeta}\cdot z\right|$ and set D its constant in the quasi triangular inequality, i.e.
$\forall a, b \in \partial\mathbb{B},\ \delta(a,b) \leq D(\delta(a,c) + \delta(c,b)).$

Lemma 3.1. *The kernel $K_t(\zeta, z) := \dfrac{t^{n(p-1)}}{\left|1-(1-t)\bar{\zeta}\cdot z\right|^{np}}$ verifies for $p > 1$:*
(H2) $|K_t(\zeta,z)| \leq Ct^{\alpha}(\delta(\zeta,z)+t)^{-\alpha-n}$, with $\alpha = n(p-1) > 0$.
(H3) For any ζ, z_0, z, t such that $\delta(z_0, z) \leq \frac{1}{2D}(t + \delta(\zeta, z_0))$ we have

$$|K_t(\zeta,z) - K_t(\zeta,z_0)| \leq C\delta(z_0,z)^{1/2}(\delta(\zeta,z_0)+t)^{-n-1/2}.$$

Proof
the condition is for small t and we have, with $\alpha = n(p-1)$

$$K_t(\zeta,z) \simeq \frac{t^{n(p-1)}}{(t+\delta(\zeta,z))^{np}} = t^{\alpha}((t+\delta(\zeta,z))^{-n-\alpha},$$

and the condition (H2).

For (H3) we have

$$|K_t(\zeta,z_0)-K_t(\zeta,z)| \simeq \frac{t^{n(p-1)}}{(t+\delta(\zeta,z_0))^{np}(t+\delta(\zeta,z))^{np}}\left|(t+\delta(\zeta,z_0))^{np}-(t+\delta(\zeta,z))^{np}\right|. \tag{3.1}$$

The function $x\longrightarrow f(x):=x^{np}$ verifies

$$|f(a)-f(b)|=|b-a|\,|f'(c)|=|b-a|\,np\,|c|^{np-1},\ c\in]a,b[$$

by the mean value property, hence here we get

$$|(t+\delta(\zeta,z_0))^{np}-(t+\delta(\zeta,z))^{np}|\leq|\delta(\zeta,z_0)-\delta(\zeta,z)|\max\left((t+\delta(\zeta,z_0))^{np-1},\ (t+\delta(\zeta,z))^{np-1}\right).$$

We have

$$t^{n(p-1)}\leq\min\left((t+\delta(\zeta,z_0))^{n(p-1)},\ (t+\delta(\zeta,z))^{n(p-1)}\right),$$

because $n(p-1)\geq 0$.

Let $A:=\max\left((t+\delta(\zeta,z_0)),\ (t+\delta(\zeta,z))\right)$, $B:=\min\left((t+\delta(\zeta,z_0)),\ (t+\delta(\zeta,z))\right)$ then putting the last two inequalities in (3.1) we get

$$|K_t(\zeta,z_0)-K_t(\zeta,z)|\leq|\delta(\zeta,z_0)-\delta(\zeta,z)|\,A^{-1}B^{-n}.$$

By the quasi triangular inequality we have

$$\delta(\zeta,z_0)\leq D\left(\delta(\zeta,z)+\delta(z,z_0)\right)\Longrightarrow t+\delta(\zeta,z)\geq D^{-1}\left(t+\delta(\zeta,z_0)\right)-\delta(z_0,z),$$

because $D\geq 1$. But we have $\delta(z_0,z)\leq\frac{1}{2D}(t+\delta(\zeta,z_0))$ hence, $t+\delta(\zeta,z)\geq\dfrac{1}{2D}\left(t+\delta(\zeta,z_0)\right)$ so we always have $A^{-1}B^{-n}\leq C(t+\delta(\zeta,z_0))^{-n-1}$.

Now we shall take advantage that $d(a,b):=\sqrt{\delta(a,b)}$ is a distance on $\partial\mathbb{B}$. We have

$$|\delta(\zeta,z_0)-\delta(\zeta,z)|=\left|d^2(\zeta,z_0)-d^2(\zeta,z)\right|=|d(\zeta,z_0)-d(\zeta,z)|\,(d(\zeta,z_0)+d(\zeta,z)).$$

Because d is a distance, $|d(\zeta,z_0)-d(\zeta,z)|\leq d(z_0,z)=\delta^{1/2}(z_0,z)$. On the other hand

$$d(\zeta,z)\leq d(\zeta,z_0)+d(z_0,z)\Longrightarrow d(\zeta,z_0)+d(\zeta,z)\leq 2d(\zeta,z_0)+d(z_0,z)\leq C\sqrt{t+\delta(\zeta,z_0)},$$

again with $\delta(z_0,z)\leq\frac{1}{2D}(t+\delta(\zeta,z_0))$. So finally we get

$$|K_t(\zeta,z_0)-K_t(\zeta,z)|\leq C\delta^{1/2}(z_0,z)(t+\delta(\zeta,z_0))^{-n-1/2},$$

and the lemma. □

3.1. **Proof of the theorem.** Suppose that the sequence S is dual bounded in $H^p(\mathbb{B})$, i.e. there is a sequence $\{\rho_a\}_{a\in S}\subset H^p(\mathbb{B})$ such that

$$\forall a\in S,\ \|\rho_a\|_p\leq C,\ \langle\rho_a,\ k_{b,p'}\rangle=\delta_{ab}.$$

Now fix $s<p$ and let $\nu\in\ell^s$. We have to show that we can interpolate the sequence ν in $H^s(\mathbb{B})$.

Let q be such that $\dfrac{1}{s}=\dfrac{1}{p}+\dfrac{1}{q}$ and write $\forall a\in S,\ \nu_a=\lambda_a\mu_a$ with $\lambda_a:=\frac{\nu_a}{|\nu_a|}|\nu_a|^{s/p}$ and $\mu_a:=|\nu_a|^{s/q}$; then $\lambda\in\ell^p,\ \mu\in\ell^q$ and $\|\nu\|_s=\|\lambda\|_p\|\mu\|_q$.

Let $k_{a,q}$ the normalized reproducing kernel for a in $H^q(\mathbb{B})$ and set $\gamma_a:=\dfrac{c(a,q)c(a,s')}{c(a,p')c(a,2)^2}$which is bounded above and below by strictly positive constants independent of $a\in\mathbb{B}$. In particular we have $\exists C>0,\ \forall a\in\mathbb{B},\ \gamma_a\leq C$.

Let $h:=\sum_{a\in S}\nu_a\gamma_a\rho_a k_{a,q}=\sum_{a\in S}\gamma_a\lambda_a\rho_a\mu_a k_{a,q}$; h takes the "good" values on S :

$$\forall b\in S,\ h(b)=\sum_{a\in S}\gamma_a\nu_a\rho_a(b)k_{a,q}(b)=\gamma_b\nu_b\rho_b(b)k_{b,q}(b)=\nu_b\,\|k_b\|_{s'},$$

because

$$\gamma_b \rho_b(b) k_{b,q}(b) = \gamma_b \left\lVert k_b \right\rVert_{p'} \frac{\left\lVert k_b \right\rVert_2^2}{\left\lVert k_b \right\rVert_q} = \gamma_b c(b,p') \frac{c(b,2)^2}{c(b,q)} (1-|b|^2)^{-n/s} = \left\lVert k_b \right\rVert_{s'},$$

by use of $\displaystyle\frac{1}{s} = \frac{1}{p} + \frac{1}{q}$.

Moreover h depends linearly on ν.

It remains to estimate its $H^s(\mathbb{B})$ norm.

We have, using Hölder inequalities

$$|h| \leq C \left(\sum_{a\in S} |\lambda_a|^p \, |\rho_a|^p \right)^{1/p} \left(\sum_{a\in S} |\mu_a|^{p'} \, |k_{a,q}|^{p'} \right)^{1/p'}.$$

Let $g := \displaystyle\left(\sum_{a\in S} |\lambda_a|^p \, |\rho_a|^p \right)^{1/p}$, we have

$$\left\lVert g \right\rVert_p^p = \int_{\partial\mathbb{B}} \sum_{a\in S} |\lambda_a|^p \, |\rho_a|^p \, d\sigma = \sum_{a\in S} |\lambda_a|^p \left\lVert \rho_a \right\rVert_p^p \leq C \left\lVert \lambda \right\rVert_p^p,$$

hence $g \in L^p(\sigma)$.

So to have h in $L^s(\sigma)$ it suffices to prove that

$$\left(\sum_{a\in S} |\mu_a|^{p'} \, |k_{a,q}|^{p'} \right)^{1/p'} \in L^q(\mathbb{B}).$$

So let $f := \displaystyle\sum_{a\in S} |\mu_a|^{p'} \, |k_{a,q}|^{p'}$ we shall show that $f \in L^{q/p'}$ $(q > p' \Longrightarrow \frac{q}{p'} > 1)$.

From $k_{a,q} = \dfrac{(1-|a|^2)^{n/q'}}{(1-\overline{a}\cdot z)^n}$, we get

$$|k_{a,q}|^{p'} = \frac{(1-|a|^2)^{\frac{np'}{q'}}}{|1-\overline{a}\cdot z|^{np'}} = (1-|a|^2)^{-\frac{np'}{q}} \frac{(1-|a|^2)^{n(p'-1)}}{|1-\overline{a}\cdot z|^{np'}} (1-|a|^2)^n.$$

Now the measure $\chi := \displaystyle\sum_{a\in S} (1-|a|^2)^n \delta_a$ is Carleson by lemma 2.2 and the kernel

$$K(a,z) := \frac{(1-|a|^2)^{n(p'-1)}}{|1-\overline{a}\cdot z|^{np'}}$$

verifies (H2) and (H3) by lemma 3.1, then we can apply the results in [3] : if χ is a Carleson measure and $\alpha \in L^r(\chi)$ then the balayage of the measure $\alpha\, d\chi$ by the kernel $K(a,z)$ is in the space $L^r(\sigma)$.

Here, let $\eta_a(z)$ be a smooth function with support in the hyperbolic ball centered at a and with radius $r > 0$ so small that these balls are disjoint for $a \in S$ and such that $\eta_a(a) = 1$. Let

$$\alpha := \sum_{a\in S} |\mu_a|^{p'} (1-|a|^2)^{-\frac{np'}{q}} \eta_a(z),$$

if $\alpha \in L^{q/p'}(\chi)$, we have that its balayage, which is precisely f, is in $L^{q/p'}(\sigma)$.

But we have

$$\int |\alpha|^{q/p'} \, d\chi = \sum_{a\in S} (1-|a|^2)^n \, |\mu_a|^q \, (1-|a|^2)^{-n} = \sum_{a\in S} |\mu_a|^q < \infty,$$

and we are done. $\square$

References

[1] E. Amar. Interpolation of interpolating sequences. *(To appear in) Indagationes Mathematicae*, 2006.
[2] E. Amar. On linear extension for interpolating sequence. In F. Meylan, editor, *Journes Complexes du Sud*, 2006.
[3] E. Amar and A. Bonami. Mesure de Carleson d'ordre α et solution au bord de l'équation $\bar{\partial}$. *Bull. Soc. Math. France*, 107:23–48, 1979.
[4] L. Carleson. An interpolation problem for bounded analytic functions. *Amer. J. Math.*, 1958.
[5] L. Hormander. A L^p estimates for (pluri-) subharmonic functions. *Math. Scand.*, 20:65–78, 1967.
[6] H. Shapiro and A. L. Shields. On some interpolation problems for analytic functions. *Amer. J. Math.*, 83:513–522, 1961.
[7] P. J. Thomas. Hardy space interpolation in the unit ball. *Indagationes Mathematicae*, 90(3):325–351, 1987.

Université Bordeaux I, 351 Cours de la Libération, 33405 Talence France.
E-mail address: `Eric.Amar@math.u-bordeaux1.fr`